

Visit

FreeTestPaper.com

for more papers



TANJONG KATONG GIRLS' SCHOOL

PRELIMINARY EXAMINATION 2016
SECONDARY FOUR

4047/01

ADDITIONAL MATHEMATICS
PAPER 1

Thursday

11 August 2016

2 h

Additional Materials: Answer Paper
Graph Paper

READ THESE INSTRUCTIONS FIRST

Write your name, class and register number on all the work you hand in.
Write in dark blue or black pen on both sides of the paper, and use a pencil for drawing graphs and diagrams.
Do not use staples, highlighters or correction fluid.

Answer **all** the questions.

Write your answers on the separate writing paper provided.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 80.

Setter : Ms Yeo
Markers : Mrs Pang / Mrs M Loy / Mdm Tan SE / Ms Yeo

This Question Paper consists of 7 printed pages, including this page.

Answer all questions

- 1 It is given that $\cos A = -\frac{1}{3}$ and $\sin B = \sqrt{\frac{2}{11}}$. A and B are in the same quadrant.
Without using a calculator, find the exact value of $\cot(90^\circ - A - B)$. [5]
- 2 (i) Find the range of values of p for which $(x+1)(x-2) > p(x+2)$ for all real values of x . [4]
(ii) Deduce the number of points at which the line $y = p(x+2)$ intersects the curve $y = (x+1)(x-2)$ for $-1 \leq p < 2$. [1]
- 3 2000 cm³ of water is transferred from a rectangular tank to an empty inverted right circular cone in 10 seconds. The ratio of the radius of the cone to the height of the cone is 1 : 3.
Find the rate of change of the horizontal surface area, A cm², of the water in the cone, when the height, h cm, of the water in the cone is 12 cm. [6]
- 4 (i) Write down and simplify, the first 3 terms in the expansion of $(2-p)^7$ in ascending powers of p . [3]
(ii) Find the value of n where n is a positive integer, given that the coefficient of x^2 is 96 in the expansion of $(1+x)^n(2-x+x^2)^7$. [4]

- 5 A curve $y = f(x)$ is such that $f''(x) = 48\sin 4x - 8\cos 2x$. The curve intersects the x -axis at P . The x -coordinate of P is $\frac{\pi}{4}$ and the gradient of the curve at P is 8. Show that $f''(x) + 16f(x) = 24\cos 2x$. [7]

- 6 The table shows experimental values of two variables x and y .

| | | | | | |
|-----|------|------|------|------|------|
| x | 2 | 4 | 6 | 7 | 8 |
| y | 1.33 | 2.29 | 3.27 | 3.77 | 6.12 |

It is known that x and y are related by an equation of the form $x^2 + \frac{y}{a} = bxy$, where a and b are constants. An error was made in recording one of the values of y .

- (i) Using a scale of 2 cm to represent 1 unit on the horizontal axis and 1 cm to represent 1 unit on the vertical axis, draw a straight line graph for the above given data. The straight line graph is to be drawn with variable x on the horizontal axis. [3]
- (ii) Use the graph to estimate
- (a) the correct value of y , [2]
- (b) the values of a and b . [2]

- (i) Express $\frac{4}{(x-3)x^2}$ in partial fractions. [4]

- (ii) Hence evaluate $\int_4^7 \frac{1}{(x-3)x^2} dx$. [4]

- 8 (i) Prove that $\frac{1-\sin x}{\cos x} + \frac{\cos x}{1-\sin x} = 2\sec x$. [3]

- (ii) In the equation

$$\frac{1-\sin x}{\cos x} + \frac{\cos x}{1-\sin x} + \tan^2 x = 2,$$

$\cos x = a$ or b where a and b are constants, and $b < 0$.

- (a) Find the value of a and of b . [2]
- (b) Solve the equation $\cos x = b$ for $-\pi \leq x \leq 2\pi$. [3]


- 9 The equation of a curve is $y = x \ln(2x - 3)$ where $x > \frac{3}{2}$.

- (i) Find the equation of the normal to the curve at $x = 2$. [4]

The normal to the curve $y = x \ln(2x - 3)$ passes through the vertex of the graph of $y = k - 4|2x + 1|$ where k is a constant.

- (ii) Determine the value of k . [2]
- (iii) Sketch the graph of $y = k - 4|2x + 1|$ for the value of k in part (ii). Show the vertex and intercepts clearly. [2]

10 Solutions to this question by accurate drawing will not be accepted.



The diagram shows a quadrilateral $HIJK$. H is the reflection of point $G(5, 7)$ in the line $x = 1$. Point $K(a, 11)$ is such that the product of the gradients of HK and JK is -3 . The perpendicular bisector of HJ intersects the x -axis at I .

(i) Deduce the coordinates of H . [1]


Find

(ii) the value of a given that $a < 0$, [2]

(iii) the equation of the perpendicular bisector of HJ , [3]

(iv) the area of quadrilateral $HIJK$. [3]

11



The diagram shows a capsule shaped object with surface area $18\pi \text{ cm}^2$. It comprised of 2 solid hemispheres of radius $r \text{ cm}$ joined to the 2 ends of a solid cylinder of radius $r \text{ cm}$ and height $h \text{ cm}$.

(i) Show that the volume, $V \text{ cm}^3$, of the object is given by $V = 9\pi r - \frac{2}{3}\pi r^3$. [4]

(ii) Find the stationary value of V , and determine if this stationary value is a maximum or minimum. [6]

THE END

Answer Key to TKGS Prelim 2016 Additional Mathematics Paper 1

| | | | |
|----------|---|---------|---|
| 1 | $7\sqrt{2}$ | 8(i) | Proof |
| | | (ii)(a) | $a=1$ and $b=-\frac{1}{3}$ |
| 2(i) | $-9 < p < -1$ | (ii)(b) | -1.91, 1.91, 4.37 |
| 2 (ii) | 1 or 2 points | | |
| | | 9(i) | $4y = -x + 2$ |
| 3 | $33\frac{1}{3} \text{ cm}^3/\text{s}$ | (ii) | $\frac{5}{8}$ |
| 4(i) | $128 - 448p + 672p^2 +$ | (iii) | |
| 4(ii) | 4 | | |
| | | | |
| 5 | proof | | |
| | | 10(i) | $(-3, 7)$ |
| 6(ii)(a) | 4.24 | (ii) | -1 |
| (b) | $a = 1, b = 2$ | (iii) | $y = 3x + 6$ |
| | | (iv) | 34 square units |
| | | | |
| 7(i) | $\frac{4}{(x-3)x^2} = \frac{4}{9(x-3)} - \frac{4}{9x} + \frac{4}{3x^2}$ | 11(ii) | 40.0 cm^3 , Stationary value of V is a maximum. |



TANJONG KATONG GIRLS' SCHOOL
PRELIMINARY EXAMINATION 2016
SECONDARY FOUR

4047/02

ADDITIONAL MATHEMATICS
PAPER 2

Friday

5 August 2016

2 h 30 min

Additional Materials: Answer Paper

READ THESE INSTRUCTIONS FIRST

Write your name, class and register number on all the work you hand in.
 Write in dark blue or black pen on both sides of the paper, and use a pencil for for any diagrams or graphs.
 Do not use staples, highlighters or correction fluid.

Answer **all** the questions.

Write your answers on the separate writing paper provided.
 Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an approved scientific calculator is expected, where appropriate.
 You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
 The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 100.

Setter : Mrs M Loy
 Markers: Mdm Tan SE, Mrs H Pang, Miss Yeo LS, Mrs M Loy

This Question Paper consists of 7 printed pages, including this page.


Answer **all** the questions

1. A man buys a new car. The value of the car depreciates with time so that its value, \$ V , after t months' use is given by $V = 132\,000e^{-pt}$, where p is a constant.
 The value of the car is expected to be \$122 000 after eight months' use.
- (i) Find the value of the car, V when the man bought it. [1]
- (ii) Show that $p = 0.01$. [2]
- (iii) Using the value of $p = 0.01$, determine the age of the car to the nearest month, when its value reached half of the original value when the man bought it. [2]
2. The function $f(x) = 1 + 2x + Ax^2 - x^3$, where A is a constant, leaves a remainder of $1\frac{3}{8}$ when divided by $(2x - 1)$.
- (i) Find the value of A . [2]
- (ii) Hence solve the equation $f(x) = 0$, giving your answers in the exact form. [2]
3. (a) (i) Solve $\sqrt{3x+2} - 3x = 0$. [2]
- (ii) On the same axes, sketch the graphs of $y = \sqrt{3x+2}$ and $y = 3x$.
 Indicate clearly all the points of intersections. [2]
- (b) The vertical height of a triangle is $\frac{8}{3-\sqrt{5}}$ cm.
 Given that the area of the triangle is $\frac{20}{\sqrt{5}-1}$ cm², without using a calculator, find the length of the base of the triangle in the form $a + b\sqrt{5}$. [3]

4. The roots of the quadratic equation, $2x^2 + 4x + 5 = 0$ are $(\alpha + 1)$ and $(\beta + 1)$.
- (i) Show that $\alpha + \beta = -4$ and hence find $\alpha\beta$. [3]
- (ii) Find the quadratic equation in x with integer coefficients, whose roots are $\frac{1}{\alpha^3}$ and $\frac{1}{\beta^3}$. [5]
5. (a) Given that $\log_2(2x + 1) - \log_4(3 - x^2) = 1$, form a quadratic equation in x and explain with clear working why the roots of the quadratic equation are real and distinct. [5]
- (b) Solve $3^{y+2} = 2(3^{-y}) + 17$. [4]

6. The curve $y = \frac{2x^2}{x^2 + 1}$ has one stationary point (p, q) .
- (i) Find the value of p and of q . [4]
- (ii) Determine whether y is increasing or decreasing for
- (a) $x > p$, [1]
- (b) $x < p$. [1]
- Hence state the nature of the stationary point. [1]
- (iii) Find $\frac{d^2y}{dx^2}$ at the stationary point and explain how $\frac{d^2y}{dx^2}$ further supports your answer in part (ii). [2]

7.



In the figure, AB is a diameter of the circle with centre O . Chords AD and BC intersect at F . AD produced meets the tangent to the circle, TBE at E . AE is an angle bisector of angle BAC .

- (i) Prove that $\angle CBD = \angle DBE$. [3]
- Given that $\angle AOF = 90^\circ$, prove that
- (ii) triangle AOF is similar to triangle ADB . [2]
- (iii) $2(AO)^2 = AF \times (AF + FD)$. [3]
8. A particle moving in a straight line passes through a fixed point O with a speed of 20 m/s. The acceleration, a m/s², of the particle, t s after passing through O is given by $a = -100e^{-3t}$. The particle comes to instantaneous rest at point N .
- (i) Find the time the particle comes to instantaneous rest at point N . [5]
- (ii) Calculate the distance ON . [4]
- (iii) Show that the average speed of the particle in the first 2 seconds rounded off to a whole number is 10 m/s. [3]

9. (i) Solve the equation $2\sin 2P = 3\cos P$ for $0^\circ \leq P \leq 360^\circ$. [4]

(ii) On the same axes, sketch for $0^\circ \leq x \leq 720^\circ$, the graphs of

$$y = \sin x \quad \text{and} \quad y = \frac{3}{2} \cos\left(\frac{x}{2}\right). \quad [4]$$

(iii) Using the solutions to part (i), determine the x -coordinates of the points of intersection of the graphs of part (ii). [4]

10. A circle, C_1 , has equation $x^2 + y^2 - 14x + 2y = -46$.

(i) Find the coordinates of the centre of the circle and the radius. [3]


The coordinates of the centre of a second circle, C_2 , is $(-4, -2)$. The equation of the tangent to the circle, C_2 at a point P is $2y = -2x + 3$.

(ii) Find the coordinates of point P . [4]

(iii) Find the exact value of the radius of C_2 and the equation of the circle, C_2 . [3]

(iv) Determine whether circles C_1 and C_2 will meet each other, showing your working clearly. [2]

11. (a) Show that $\frac{d}{dx}(2x\sqrt{2x+3}) = \frac{6x+6}{\sqrt{2x+3}}$ [3]




The diagram shows part of the curve $y = \frac{3x+3}{\sqrt{2x+3}}$. The curve intersects the x -axis at point A . The line through A and perpendicular to the line, $y + x = -7$ intersects the curve again at another point, B .

(i) Show that the y -coordinate of point B is 4. [5]

(ii) Given that the line AB intersects the y -axis at C , determine the area of the shaded region bounded by the line CB , the curve, the line $x = 5$, the x -axis and the y -axis. [4]

End of Paper

TKGS S4 PRELIM 2016 Answer Key:

| | | | |
|--------|--|---------|--|
| 1(i) | $V = 132\,000$ | (ii) | show |
| (iii) | 70 months | | |
| 2(i) | $A = -2$ | (ii) | $x = 1, \frac{-3 \pm \sqrt{5}}{2}$ |
| 3(a)i | $x = \frac{2}{3}$ | ii |  |
| (b) | $\frac{5\sqrt{5}}{2} - \frac{5}{2}$ | 4(i) | $\alpha\beta = \frac{11}{2}$ |
| i(ii) | $1331x^2 - 16x + 8 = 0$ | 5(a) | Discriminant = 368 Since discriminant > 0 , the roots of the quadratic equation are real and distinct. |
| 5(b) | $y = 0.631$ | 6(i) | $p = 0, q = 0$ |
| (ii)a | $\frac{dy}{dx} > 0, y$ is increasing | 6(ii)b | $\frac{dy}{dx} < 0, y$ is decreasing |
| | Since the value of $\frac{dy}{dx}$ changes from negative to positive value, the stationary point is a minimum point. | 6(iii) | $\frac{d^2y}{dx^2} = 4$, since $\frac{d^2y}{dx^2} > 0$, the stationary point is minimum, thus reiterating the result from part (ii). |
| | proof | 8(i) | $t = 0.305$ s |
| (ii) | Distance = 2.59 m | 8(iii) | show |
| 9(i) | $48.6^\circ, 90^\circ, 131.4^\circ, 270^\circ$ | (ii) | |
| 9(iii) | $97.2^\circ, 180^\circ, 262.8^\circ, 540^\circ$ | 10(i) | Centre(7, -1), radius = 2 units |
| 10(ii) | $P(-\frac{1}{4}, \frac{7}{4})$ | 10(iii) | Radius = $\frac{15\sqrt{2}}{4}$ units |

| | | | |
|--------|--|---------|--|
| | | | $(x+4)^2 + (y+2)^2 = (\frac{15\sqrt{2}}{4})^2 / \frac{225}{8}$ |
| (iv) | Sum of radii(7.30 units) < distance between the centres (11.0 units) thus the circles will not meet. | 11(a) | show |
| 11(b)i | show | 11(b)ii | 16.5 units ² |