

*Visit*

**FREETESTPAPER.com**

*for more papers*




Website: [freetestpaper.com](http://www.freetestpaper.com)



[Facebook.com/freetestpaper](https://www.facebook.com/freetestpaper)



[Twitter.com/freetestpaper](https://www.twitter.com/freetestpaper)



**ANDERSON SECONDARY SCHOOL**  
**Preliminary Examination 2020**  
**Secondary Four Express & Five Normal**



CANDIDATE NAME:

CLASS:

INDEX NUMBER:

---

**ADDITIONAL MATHEMATICS**

**4047/01**

Paper 1

**4 August 2020**

**2 hours**

**0800 – 1000h**

Candidates answer on the Question Paper.

No Additional Materials are required.

---

**READ THESE INSTRUCTIONS FIRST**

Write your name, class and index number in the spaces at the top of this page.

Write in dark blue or black pen.

You may use HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid/tape.

Answer **all** the questions.

Omission of essential working will result in loss of marks.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an approved scientific calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the test, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The total number of marks for this paper is 80.

**Mathematical Formulae****1. ALGEBRA****Quadratic Equation**

For the equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

**Binomial expansion**

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \dots + \binom{n}{r} a^{n-r} b^r + \dots + b^n,$$

where  $n$  is a positive integer and  $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1) \dots (n-r+1)}{r!}$

**2. TRIGONOMETRY****Identities**

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

**Formulae for  $\Delta ABC$** 

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} ab \sin C$$

**1** Given that  $\theta$  is obtuse and  $\tan \theta = a$ , express, in terms of  $a$ ,

**(i)**  $\cos \theta$ , [3]

**(ii)**  $\operatorname{cosec} \theta$ . [2]

- 2 Find the set of values of the constant  $k$  for which the curve  $y = -x^2 + (1-k)x - 2$  lies entirely below the line  $x + y = 0$ . [4]

- 3 Given that  $y = he^x + \frac{k}{e^{2x}}$ , and that  $\frac{d^2y}{dx^2} - 2\frac{dy}{dx} = e^x + 2e^{-2x}$ , find the value of each of the constants  $h$  and  $k$ . [4]

- 4 A cylindrical ice block of base radius  $r$  cm is melting in such a way that the total surface area,  $A$  cm<sup>2</sup>, is decreasing at a constant rate of  $72$  cm<sup>2</sup>/s. Given that the height is twice the radius and assuming that the ice block retains its shape, calculate the rate of change of  $r$  when  $r = 5$ . [4]

- 5** Ms Lee bought her dream car last year. The value, \$ $A$ , of the car is given by the formula  $A = 150000e^{-pt}$ , where  $p$  is a constant and  $t$  is the age of the car in months. The value of the car after 2 years is expected to be \$120 000.
- (i) Find the amount which Ms Lee paid for the car. [1]
- (ii) Determine the value of the car after 40 months. Give your answer to the nearest dollar. [3]
- (iii) Find the age of the car when its value drops to \$60 000. Give your answer to the nearest month. [2]

6 (i) Factorise  $2x^3 + 3x^2 - 8x - 12$ . [3]

(ii) Hence solve the equation  $\frac{x^3}{4} + \frac{3}{4}x^2 - 4x - 12 = 0$ . [3]

- 7 (i) Write down and simplify the first three terms in the expansion, in descending powers of  $x$ , of  $\left(1 - \frac{2}{x}\right)^8$ . [2]

- (ii) Given that there is no  $x$  term in the expansion of  $(1 - 2x - kx^2)\left(1 - \frac{2}{x}\right)^8$ , find the constant term in the expansion. [4]

8 The equation of a curve is  $y = 2 - x - \frac{2x+3}{x-3}$ .


(i) Find  $\frac{dy}{dx}$  and  $\frac{d^2y}{dx^2}$ .

[3]

(ii) Find the  $x$ -coordinate(s) of the stationary point(s) of the curve. [3]

(iii) Determine the nature of each stationary point. [2]

9



The diagram shows a quadrilateral  $ABCD$  in which  $AB = BC$ ,  $\angle BCD = 90^\circ$  and  $D$  is a point on the  $y$ -axis. The coordinates of the points  $A$ ,  $B$  and  $C$  are  $(3, 2)$ ,  $(8, 2)$  and  $(11, k)$  respectively.

- (i) Given that  $k > 2$ , show that  $k = 6$ . [3]

(ii) Find the coordinates of  $D$ . [3]


(iii) Find the length of  $CD$  and hence find the area of the quadrilateral  $ABCD$ . [5]

- 10 (a) Find the values of  $x$  and  $y$  which satisfy the equations

$$\frac{1}{\sqrt{e^{2x-4y}}} = \frac{\sqrt[3]{e}}{e},$$
$$\frac{10^y}{2^x} = 2(5^{x+1}).$$

[5]

(b)



A solid triangular pyramid  $ABCD$ , with base  $ABC$  and vertex  $D$  such that  $D$  is vertically above  $A$ , has a base area of  $(8 + 2\sqrt{5}) \text{ cm}^2$  and height  $(12 - \sqrt{5}) \text{ cm}$ .

The top part of the pyramid is removed by a cut parallel to its base and passing through the midpoint of  $AD$ . Find the volume of the remaining solid, leaving

your answer in the form  $\frac{7(a + b\sqrt{5})}{12} \text{ cm}^3$ , where  $a$  and  $b$  are integers. [4]

- 11** A particle moving in a straight line passes a fixed point  $O$  with a velocity  $6 \text{ ms}^{-1}$ .  
The acceleration of the particle,  $a \text{ ms}^{-2}$ , is given by  $a = 2t - 5$ , where  $t$  seconds is the time after passing  $O$ . Find
- (i) the values of  $t$  when the particle is instantaneously at rest, [3]

(ii) the displacement of the particle from  $O$  at  $t = 3$ , [3]

(iii) the total distance travelled by the particle in the first 3 seconds of its motion. [2]

**12** It is given that  $f(x) = 2\sin\frac{x}{2}$  and  $g(x) = 3\cos x + 1$  where  $0 \leq x \leq 2\pi$ .

**(i)** State the period of  $f(x)$ . [1]

**(ii)** State the smallest value of  $f(x)$ . [1]


**(iii)** State the largest value of  $g(x)$ . [1]

**(iv)** State the largest value of  $|f(x) - g(x)|$ . [1]

- (v) Sketch, on the same axes, the graphs of  $y = f(x)$  and  $y = g(x)$  for  $0 \leq x \leq 2\pi$ . [4]

- (vi) Given that the solutions to the equation  $f(x) = g(x)$  for  $0 \leq x \leq 2\pi$  are  $a$  and  $b$  where  $a < b$ , state the range of value of  $x$  for which  $f(x) \geq g(x)$ . [1]

End of Paper



**ANDERSON SECONDARY SCHOOL**  
**Preliminary Examination 2020**  
**Secondary Four Express & Five Normal**



CANDIDATE NAME:

CLASS:

INDEX NUMBER:

**ADDITIONAL MATHEMATICS**

**4047/02**

Paper 2

**11 August 2020**

**2 hours 30 minutes**

**0800 – 1030h**

Candidates answer on the Question Paper.

No Additional Materials are required.

**READ THESE INSTRUCTIONS FIRST**

Write your class, index number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid/tape.

Answer **all** the questions.

Omission of essential working will result in loss of marks.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an approved scientific calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The total number of marks for this paper is 100.

This document consists of **22** printed pages.

Setter: Mdm Mirshasha

**Mathematical Formulae****1. ALGEBRA****Quadratic Equation**

For the equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

**Binomial expansion**

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \dots + \binom{n}{r} a^{n-r} b^r + \dots + b^n,$$

where  $n$  is a positive integer and  $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\dots(n-r+1)}{r!}$

**2. TRIGONOMETRY****Identities**

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

**Formulae for  $\Delta ABC$** 

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} ab \sin C$$

1 The roots of the quadratic equation  $5x - 2x^2 = 2 - k$  are  $\alpha$  and  $\beta$ . The roots differ by  $4\frac{1}{2}$ .

(i) Show that  $k = 9$ . [4]

(ii) Hence find a quadratic equation with integer coefficients whose roots are  $\alpha^3$  and  $\beta^3$ . [4]

- 2 (a) It is given that  $\tan(A+B) = 8$  and  $\tan B = 2$ . **Without using a calculator**, find the exact value of  $\cot A$ . [3]

- (b) (i) Prove that  $\sin 2x(\cot x - \tan x) = 2 \cos 2x$ . [3]

(ii) Hence solve the equation  $\sin 2x(\cot x - \tan x) = \sec 2x$  for  $0 \leq x \leq \pi$ .

[4]

3 The equation of circle  $C_1$ , with centre  $A$ , is  $x^2 + y^2 - 8x - 4y + 11 = 0$ .

(i) Find the coordinates of  $A$  and the radius of  $C_1$ . [3]

(ii) Explain why the line  $x = 1$  is a tangent to  $C_1$ . [1]

The equation of a second circle  $C_2$ , with centre  $B$ , is  $(x+3)^2 + (y-2)^2 = R^2$ .


(iii) Find the length of  $AB$ . [2]

(iv) State the possible value(s) of  $R$  such that  $C_1$  and  $C_2$  touch each other at exactly one point. [2]

- 4 (a) Given that  $\log_{\sqrt{2}} x = m$  and  $\log_4 y = n$ , express  $\frac{\sqrt{y}}{x^6}$  in terms of  $m$  and  $n$ . [4]

(b) Solve the equation  $\log_2(3x-2) = \log_4(x^2+1) + \frac{2}{\log_{\sqrt{2}} 2}$ . [6]

- 5 The diagram shows a solid cylinder of radius  $r$  cm and height  $h$  cm inscribed in a hollow cone of height 45 cm and base radius 20 cm. The cylinder rests on the base of the cone and the circumference of the top surface of the cylinder touches the curved surface of the cone.



- (i) Show that the volume,  $V$  cm<sup>3</sup>, of the cylinder is given by  $V = 45\pi r^2 - \frac{9}{4}\pi r^3$ . [3]

- (ii) Given that  $r$  can vary, find the maximum volume of the cylinder, leaving your answer in terms of  $\pi$ . [4]


- (iii) Hence show that the cylinder occupies at most  $\frac{4}{9}$  of the volume of the cone. [2]

- 6 (i) Express  $\frac{2x^3 - 20x^2 - 17x - 10}{(x^2 - 4)(2x^2 + 1)}$  in partial fractions. [5]

(ii) Differentiate  $\ln(2x^2 + 1)$  with respect to  $x$ . [2]

(iii) Using the results from parts (i) and (ii), find  $\int \frac{2x^3 - 20x^2 - 17x - 10}{(x^2 - 4)(2x^2 + 1)} dx$ . [3]

7



The diagram shows the curve  $y = f(x)$  which intersects the  $x$ -axis at points  $A$  and  $B$ .

The tangent to the curve at the point  $\left(\frac{1}{2}, 10\right)$  intersects the  $x$ -axis at point  $C$ . It is

given that  $\frac{dy}{dx} = -8x + 10$ .

- (i) Show that  $B$  is the point  $(3, 0)$  and find the coordinates of  $C$ . [5]

(ii) Find the area of the shaded region.

[5]


8 (i) Sketch the graph of  $y = |x^2 - 17x + 16|$ . [3]

(ii) Find the range of values of  $x$  for which  $|x^2 - 17x + 16| > 54$ . [5]

- (iii) Given that  $|x^2 - 17x + 16| = k$  has more than 2 distinct solutions, state the range of values of  $k$ . [2]

- (iv) Determine the number of solutions of the equation  $|x^2 - 17x + 16| = 2x - 2$ . Justify your answer. [2]

- 9 The diagram shows the side view  $PQRST$  of a tent. The tent rests with  $RS$  on horizontal ground.  $PQRST$  is symmetrical about the vertical  $PU$ , where  $U$  is the midpoint of  $RS$ . Angle  $QPU = \text{angle } QRU = \theta$  radians and the lengths of  $PQ$  and  $QR$  are 90 cm and 150 cm respectively. The vertical height of  $P$  from the ground is  $h$  cm.



- (i) Explain clearly why  $h = 90 \cos \theta + 150 \sin \theta$ . [2]

- (ii) Express  $h$  in the form  $R \cos(\theta - \alpha)$ , where  $R > 0$  and  $\alpha$  is an acute angle. [4]


- (iii) Find the greatest possible value of  $h$  and the value of  $\theta$  at which this occurs. [3]

- (iv) Find the values of  $\theta$  when  $h = 160$ . [3]

- 10** A bowl of liquid is heated to a high temperature. It subsequently cools in such a way that its temperature,  $T^{\circ}\text{C}$ , is given by  $T = 15 + Ae^{-kt}$ , where  $t$  minutes is the time of cooling and  $A$  and  $k$  are constants. The table below shows corresponding values of  $t$  and  $T$ .

$t$	5	10	15	20	25
$T$	58.8	40.3	29.6	23.4	19.9

- (i) Draw the graph of  $\ln(T - 15)$  against  $t$ . [3]



(ii) Use the graph to estimate the value of each of the constants  $A$  and  $k$ . [5]

(iii) State the initial temperature of the liquid. [1]

(iv) Use the graph to estimate the time taken for the temperature of the liquid to drop to half of its original temperature. [2]

**End of Paper**

